


# Dune-slope Activity due to Frost and Wind, Throughout the North Polar Erg, Mars




S. Diniega<sup>1</sup>, C.J. Hansen<sup>2</sup>, A. Allen<sup>3</sup>, N. Grigsby<sup>4</sup>, Z. Li<sup>5</sup>, T. Perez<sup>6</sup>, M. Chojnacki<sup>7</sup>.

<sup>1</sup>Jet Propulsion Laboratory, California Institute of Technology (4800 Oak Grove, Pasadena, CA 91109; [serina.diniega@jpl.nasa.gov](mailto:serina.diniega@jpl.nasa.gov)), <sup>2</sup>Planetary Science Institute, <sup>3</sup>Santa Barbara City College, <sup>4</sup>Boise State University, <sup>5</sup>Oxford University, <sup>6</sup>California Institute of Technology, <sup>7</sup>Lunar and Planetary Laboratory, University of Arizona.




Hansen et al. (2011) noted that sublimation activity appeared enhanced within new alcoves (identified in the subsequent summer/defrosted images) and thus proposed a connection (as has been proposed for southern dune-gullies (e.g., Diniega et al., 2010; Dundas et al., 2012; 2015)).

While it appears that alcoves and aprons do grow during the spring, initial alcove formation occurs early enough for a layer of frost to be deposited over the feature.



*During the summer,  
Alcoves fill-in,  
fade away ...  
But no new ones  
form.*



Hansen et al. (2015) confirmed the appearance of alcoves under the frost within a couple of fields, but constrained activity to fall/winter. Our study greatly expanded the fields and Mars years examined, and constrains the formation mechanism.

We hypothesize that the alcove formation occurs during the autumn-early winter period and is initiated, in some way, by early frost processes.

Further constraints: alcoves form regularly over all areas of the steep (downwind) slopes. Also, alcove mass-wasting is active only through one Mars year's seasonal cycle (i.e., they do not re-activate).



**After much data:  
We think autumn frost moves sand,  
Summer winds erase.**

Via the rate of alcove erasure (a proxy for wind-driven sand transport), we estimate an "effective" aeolian sand fluxes of 1-few m<sup>3</sup>/Mars year (similar to Bridges et al., 2013).

Comparing this to the rate of alcove formation, we find that alcove-formation-driven transport of sand makes up about 1/5-1/50<sup>th</sup> of the total sand flux.

Thus: both aeolian sand transport and alcove formation processes are significantly modifying the dunes within the martian north polar erg, and both wind-driven and seasonal-frost driven processes should be considered (together) within future interpretations of polar dune morphology and evolution

**So what moves the dunes?  
Need to consider alcoves  
with standard dune winds.**



**Acknowledgements:**  
This research supported by MDAP, JPL summer interns, HiRISE and

SD's work was carried out at JPL under a contract with NASA. The full study was supported by MDAP grant NNN13D465T, and MC is also supported by MDAP grant NNN14ZDA001N.

**References:** Bridges et al. 2013, Bedform migration on Mars: Current results and future plans. *Aeolian Res.* 9, 133-151, doi:10.1016/j.aeolia.2013.02.004. • Diniega et al., 2010. Seasonality of present-day Martian dune-gully activity. *Geology* 38, 1047-1050, doi:10.1130/G31287.1. • Diniega et al., submitted. Dune-slope activity due to frost and wind, throughout the north polar erg, Mars. *Geologic Society London (special issue Martian Gullies & their Earth Analogues)*. • Dundas et al. 2012. Seasonal activity and morphological changes in Martian gullies. *Icarus* 220, 124-143, doi:10.1016/j.icarus.2012.04.005. • Dundas et al. 2015. Long-term monitoring of martian gully formation and evolution with MRO/HiRISE. *Icarus* 251, 244-263, doi:10.1016/j.icarus.2014.05.013. • Hansen et al. 2011. Seasonal erosion and restoration of Mars' northern polar dunes. *Science* 331, 575-578, doi:10.1126/science.1197636. • Hansen et al. 2015. Agents of change on Mars' northern dunes: CO<sub>2</sub> ice and wind. *Icarus* 251, 264-274, doi:10.1016/j.icarus.2014.11.015. • Horgan & Bell 2012. Seasonally active slipface avalanches in the north polar sand sea of Mars: Evidence for a wind-related origin. *Geophys. Res. Lett.* 39, L09201, doi:10.1029/2012GL051329.